



MAASAI MARA UNIVERSITY

MAIN EXAMINATION 2018/2019 ACADEMIC YEAR SECOND YEAR SECOND SEMESTER EXAMINATIONS

FOR

THE DEGREE OF BACHELOR OF SCIENCE

MAT 2213: CLASSICAL MECHANICS

DATE :

TIME:

DURATION: 2HRS

INSTRUCTIONS TO CANDIDATES

- 1. This paper contains FOUR (4) questions**
- 2. Answer question ONE (1) and any other TWO (2) questions**
- 3. Do not forget to write your Registration Number.**

QUESTION ONE (30 MARKS)

- a) Show that the acceleration \mathbf{a} of a particle which travels along a space curve with velocity \mathbf{v} is given by

$$\mathbf{a} = \frac{dv}{dt} \mathbf{T} + \frac{v^2}{R} \mathbf{N} \text{ where } \mathbf{T} \text{ is the unit tangent vector to the space curve,}$$

\mathbf{N} is the unit principal normal and R is the radius of curvature. **(4mks)**

- b) Find the impulse developed by a force given by $\mathbf{F} = 4t\mathbf{i} + (6t^2 - 2)\mathbf{j} + 12\mathbf{k}$ from $t=0$ to $t=2$. **(3mks)**

- c) A particle moves from rest in a circular path of a circle of radius 20 cm. If its tangential speed is 40 cm/sec, calculate its angular velocity, angular acceleration and normal acceleration. **(5mks)**

- d) The angular momentum of a particle is given as a function of time t by

$$\Omega = 6t^2\mathbf{i} - (2t + 1)\mathbf{j} + (12t^3 - 8t^2)\mathbf{k}. \text{ Find the torque at time } t=1. \quad \mathbf{(3mks)}$$

- e) An object of mass m is dropped from a height H above the ground. Prove that if air resistance is negligible, then it will reach the ground in

i. in a time $\sqrt{2H/g}$ **(2mks)**

ii. with speed $\sqrt{2gH}$ **(2mks)**

- f) Given a space curve C with position vector

$$\mathbf{r} = 3\cos 2t\mathbf{i} + 3\sin 2t\mathbf{j} + (8t - 4)\mathbf{k}. \text{ Find the}$$

i. Curvature **(4mks)**

ii. Unit principal normal to any point of the space curve **(3mks)**

- g) A particle of mass m moves along the x axis under the influence of a conservative force field having potential $V(x)$. If the particle is located at positions x_1 and x_2 at respective times t_1 and t_2 , prove that if E is the total energy,

$$t_2 - t_1 = \sqrt{\frac{m}{2}} \int_{x_1}^{x_2} \frac{dx}{\sqrt{E - V(x)}} \quad \mathbf{(4mks)}$$

QUESTION TWO (20 MARKS)

- a) At time $t=0$ a parachutist having weight of magnitude mg is located at $z=0$ and is travelling vertically downward with speed v_0 . If the force of air resistance acting on the parachute is proportional to the instantaneous speed, find the
- i. Speed **(5mks)**
 - ii. Distance travelled **(3mks)**
 - iii. Acceleration at any time $t>0$ **(2mks)**
 - iv. Show that the parachutist approaches a limiting speed given by $\frac{mg}{\beta}$, where β is an arbitrary constant. **(2mks)**
- b) Find the work done in moving a particle once around a circle C in the xy plane, if the circle has center at the origin and radius 3 and if the force field is given by
- $$\mathbf{F} = (2x - y + z)\mathbf{i} + (x + y - z^2)\mathbf{j} + (3x - 2y + 4z)\mathbf{k} \quad \textbf{(8mks)}$$

QUESTION THREE (20 MARKS)

- a) Determine the motion of a simple pendulum of length L and mass m assuming small vibrations and no resisting forces. Hence determine the period and amplitude and frequency of vibrations. **(13mks)**
- b) A cannon has its maximum range given by R_{\max} . Prove that
- i. The height reached in such a case is $\frac{1}{4}R_{\max}$ **(4mks)**
 - ii. the time of flight is $\sqrt{2R_{\max}/g}$ **(3mks)**

QUESTION FOUR (20 MARKS)

- a) i) Show that $\mathbf{F} = (2xy + z^3)\mathbf{i} + x^2\mathbf{j} + 3xz^2\mathbf{k}$ is a conservative force field. **(3mks)**
- ii) Find the scalar potential. **(4mks)**
- iii) Find the work done in moving an object in this field from (1,-2,1) to (3,1,4). **(3mks)**
-
- b) A spring of negligible mass, suspended vertically from one end, is stretched distance of 20cm when a 5g mass is attached to the other end. The spring and mass are placed on a horizontal frictionless table with the suspension point fixed. The mass is pulled away a distance 20cm beyond the equilibrium position O and released. Find
- i. The differential equation and initial conditions describing the motion. **(3mks)**
- ii. The position at any time t, **(4mks)**
- iii. The amplitude, period and frequency of the vibrations. **(3mks)**