



MAASAI MARA UNIVERSITY

**REGULAR UNIVERSITY EXAMINATIONS
2022/2023 ACADEMIC YEAR
SECOND YEAR FIRST SEMESTER**

**MASTER OF SCIENCE IN ECONOMICS
AND STATISTICS**

ECS 8202: MULTIVARIATE ANALYSIS

DATE 6TH FEBRUARY 2023

TIME: 1430HRS-1730HRS

Instructions:

1. Answer questions **one** and any other **two**.
 2. Candidates are advised not to write on the question paper.
 3. Candidates must hand in their answer booklets to the invigilator while in the examination room.
-
-

QUESTION ONE (30 Marks)

a) Given that $X \sim N_3(\mu, \Sigma)$ with

$$\mu = \begin{pmatrix} 40 \\ 28 \\ 12 \end{pmatrix} \quad \Sigma = \begin{pmatrix} 36 & -6 & -12/5 \\ -6 & 16 & 4 \\ -12/5 & 4 & 4 \end{pmatrix}$$

Find;

- i. The distribution of $Y = \begin{pmatrix} X_1 - X_3 \\ X_1 - X_2 - X_3 \end{pmatrix}$
(3 marks)
 - ii. The correlation matrix for the data
(3 marks)
 - iii. The distribution of X_1 given that $X_2=23$ and $X_3=14$.
(3 marks)
- iv. The partial correlation coefficient between X_1 and X_2 for fixed values of X_3 and that of X_1 and X_3 for the fixed values of X_2 .
(2 marks)

b) Use the data below for a bivariate normal distribution to test at $\alpha = 0.05$ level the hypothesis $H_0 = (17 \ 15)'$ vs $H_1 \neq (17 \ 15)'$.

$$X = \begin{pmatrix} 15 & 13 & 12 & 15 & 17 & 10 & 16 \\ 19 & 15 & 17 & 21 & 24 & 20 & 17 \end{pmatrix}$$

(5 marks)

c) Given that $\bar{X}_1 = (33 \ 12 \ 10)'$ comes from population I and

$\bar{X}_2 = (27 \ 19 \ 8)'$ Comes from population II and both populations have the same sample covariance matrix.

$$S = \begin{pmatrix} 20 & -4 & 15 \\ -4 & 16 & 0 \\ 15 & 0 & 4 \end{pmatrix}$$

Use discriminant rule to classify $X = (34 \ 11 \ 8)'$

(5 marks)

(c) For the matrix

$$A = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{pmatrix}$$

- i. Could A be a covariance matrix? Explain **(2 marks)**
- ii. Obtain determinant of A and A^{-1} **(3 marks)**
- iii. Compute the spectral decomposition of A **(4 marks)**

QUESTION TWO (20 Marks)

i. Observations on three responses are collected for two treatments as shown in the table below.

| Treatment | 1 | 1 | 1 | 2 | 2 |
|----------------|----|----|----|----|----|
| x ₁ | 12 | 13 | 8 | 11 | 10 |
| x ₂ | 19 | 18 | 14 | 11 | 15 |
| x ₃ | 8 | 9 | 7 | 14 | 12 |

find;

- i) The matrix of sum of squares due to the treatment. **(4 marks)**
 - ii) The matrix of residual sum of squares **(5 marks)**
- ii. Let random variables $\underline{x}' = [x_1, x_2, x_3]$ be distributed as $N_3(\underline{\mu}, \underline{\Sigma})$

$$\underline{\mu} = \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \text{ and } \Sigma = \begin{bmatrix} 4 & 1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 3 \end{bmatrix}$$

Find the following

- i.** Correlation matrix of \underline{x} **(2 marks)**
- ii.** The distribution of $z = 4x_1 - 6x_2 + x_3$ **(3 marks)**
- iii.** The distribution of $z = \begin{pmatrix} x_1 - x_2 + x_3 \\ 2x_1 + x_2 - x_3 \end{pmatrix}$ **(3 marks)**
- iv.** The Wilk's lambda statistic and use it to test the hypothesis that there is no treatment effects (use $\alpha = 0.05$) **(3 marks)**

QUESTION THREE (20 MARKS)

- a)** Describe briefly any two objectives of scientific investigation based on multivariate data **(2 marks)**
- b)** Briefly explain the idea behind MANOVA stating all the assumptions. **(2 marks)**
- c)** Distinguish between Factor Analysis and Canonical correlation Analysis **(2 marks)**
- d)** Consider data matrix for $n=3$ for a bivariate distribution

$$X = \begin{bmatrix} 6 & 10 & 8 \\ 9 & 6 & 3 \end{bmatrix}$$

$$\bar{X} = \begin{bmatrix} 8 \\ 6 \end{bmatrix}$$

i. Evaluate the observed T^2 for $\underline{\mu}'_0 = [9 \ 5]$.

(6 marks)

ii. What is the sampling distribution of T^2 in this case?

(6 marks)

QUESTION FOUR (20 MARKS)

a) Let $X \sim N(\mu, \Sigma)$ be a trivariate normal random vector. Suppose a certain

sample gave $S = \begin{pmatrix} 64 & 0 & 9.6 \\ 0 & 16 & 0 \\ 9.6 & 0 & 36 \end{pmatrix}$

find

i. The eigen values of this matrix

(4 marks)

ii. The first two principal components

(4 marks)

iii. The total variance explained by the two components.

(4 marks)

b) Consider the covariance matrix

$$\Sigma = \begin{bmatrix} 1 & 4 \\ 4 & 100 \end{bmatrix}$$

And the derived correlation matrix

$$\rho = \begin{bmatrix} 1 & 0.4 \\ 0.4 & 100 \end{bmatrix}$$

Determine the principal components for Σ providing percentage of explained variability for each variate.

(6 marks)

QUESTION FIVE (20 Marks)

a) Suppose

$$\Sigma \text{ (covariance matrix)} = \begin{bmatrix} 4 & 1 & 2 \\ 1 & 9 & -3 \\ 2 & -3 & 25 \end{bmatrix}$$

Obtain standard deviation and population correlation matrix in the form of $v^{1/2}$ and ρ respectively.

(5 marks)

b) Given the deviation vectors

$$e_1 = \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} \text{ and } e_2 = \begin{pmatrix} -2 \\ 0 \\ 2 \end{pmatrix}$$

Compute the sample variance-covariance matrix S_n and the sample correlation matrix γ using geometric concept.

(3 marks)

c) The classic blue pullovers data given below is a data set consisting of 10 measurements of 4 variables. A textile shop manager is studying the sales of classic blue pullovers over 10 periods. He uses three different marketing methods and hopes to understand his sales as a fit of these variables using statistics. The variables measured are:

| Sno. | Sales | Price | Advert | Ass.Hours |
|-------------|--------------|--------------|---------------|------------------|
| 1 | 230 | 125 | 200 | 109 |
| 2 | 181 | 99 | 55 | 107 |
| 3 | 165 | 97 | 105 | 98 |
| 4 | 150 | 115 | 85 | 71 |
| 5 | 97 | 120 | 0 | 82 |
| 6 | 192 | 100 | 150 | 103 |
| 7 | 181 | 80 | 85 | 111 |
| 8 | 189 | 90 | 120 | 93 |
| 9 | 172 | 95 | 110 | 86 |
| 10 | 170 | 125 | 130 | 78 |

Note: X_1 : Number of sold pullovers

X_2 : Price in (EUR)

X_3 : Advertisement costs in local newspapers (in EUR)

X_4 : Presence of a sales assistant (in hours per period)

i. Calculate the vector of the means for the four variables in the dataset
(4 Marks)

ii. Calculate the sample covariance matrix
(4 Marks)

iii. Calculate the sample correlation matrix
(4 Marks)

END//=